


## Colorado State University Veterinary Medical Center Feline Acute Pain Scale

Pain Score	Example	Psychological & Behavioral	Response to Palpation	Body Tension
No Score		<input type="checkbox"/> Animal is sleeping and cannot be evaluated		
0		<input type="checkbox"/> Content and quiet when unattended <input type="checkbox"/> Comfortable when resting <input type="checkbox"/> Interested in or <b>curious</b> about surroundings	<input type="checkbox"/> <b>Not bothered</b> by palpation of wound or surgery site, or to palpation elsewhere	Minimal
1		<input type="checkbox"/> Signs are often subtle and not easily detected in the hospital setting; more likely to be detected by the owner(s) at home <input type="checkbox"/> Earliest signs at home may be <u>withdrawal from surroundings or change in normal routine</u> <input type="checkbox"/> In the hospital, may be content or <b>slightly unsettled</b> <input type="checkbox"/> <b>Less interested</b> in surroundings but will look around to see what is going on	<input type="checkbox"/> May or may not react to palpation of wound or surgery site	Mild
2		<input type="checkbox"/> Decreased responsiveness, <b>seeks solitude</b> <input type="checkbox"/> <b>Quiet</b> , loss of brightness in eyes <input type="checkbox"/> <b>Lays curled up or sits tucked up</b> (all four feet under body, shoulders hunched, head held slightly lower than shoulders, tail curled tightly around body) with eyes partially or mostly closed <input type="checkbox"/> <b>Hair coat appears rough</b> or fluffed up <input type="checkbox"/> May intensively groom an area that is painful or irritating <input type="checkbox"/> Decreased appetite, <b>not interested in food</b>	<input type="checkbox"/> <b>Responds aggressively or tries to escape</b> if painful area is palpated or approached <input type="checkbox"/> Tolerates attention, may even perk up when petted as long as painful area is avoided	Mild to Moderate <b>Reassess analgesic plan</b>
3		<input type="checkbox"/> Constantly <b>yowling, growling, or hissing</b> when unattended <input type="checkbox"/> May bite or chew at wound, but <b>unlikely to move</b> if left alone	<input type="checkbox"/> <b>Growls or hisses at non-painful palpation</b> (may be experiencing allodynia, wind-up, or fearful that pain could be made worse) <input type="checkbox"/> <b>Reacts aggressively</b> to palpation, <b>adamantly pulls away</b> to avoid any contact	Moderate <b>Reassess analgesic plan</b>
4		<input type="checkbox"/> Prostrate <input type="checkbox"/> Potentially <b>unresponsive</b> to or unaware of surroundings, difficult to distract from pain <input type="checkbox"/> Receptive to care (even mean or wild cats will be more tolerant of contact)	<input type="checkbox"/> <b>May not respond</b> to palpation <input type="checkbox"/> <b>May be rigid to avoid painful movement</b>	Moderate to Severe <b>May be rigid to avoid painful movement</b> <b>Reassess analgesic plan</b>



Comments \_\_\_\_\_